# Author Index


Bacchi, C.J., see Fairlamb, A.H., 185
Bishop, R.P. and Miles, M.A.
Chromosome size polymorphisms of *Leishmania donovani*, 263
Bogitsh, B.J., see Scallon, B.J., 237
Bordier, C., see Bouvier, J., 73
Boschetti, M.A., Piras, M.M., Henriquez, D. and Piras, R.
The interaction of a *Trypanosoma cruzi* surface protein with Vero cells and its relationship with parasite adhesion, 175
Bouvier, J., Etges, R. and Bordier, C.
Identification of the promastigote surface protease in seven species of *Leishmania*, 73
Bremner, A.F., see Lockwood, B.C., 89
Brénière, S.F., see Legrand, D., 117

Campbell, T., see Komuniecki, R., 147
Capron, A., see Legrand, D., 117
Carter, C.E., see Scallon, B.J., 237
Cerami, A., see Fairlamb, A.H., 185
Cerami, A., see Henderson, G.B., 39
Chang, Y,-D., see King, D.L., 47
Christie, D.A., Powell, J.W., Stables, J.N. and Watt, R.A.
A nuclear magnetic resonance study of the role of phosphoenol pyruvate carboxykinase (PEPCK) in the glucose metabolism of *Dipetalonema viteae*, 125
Cook, G.A., see Nasir, A., 295
Coombs, G.H., see Lockwood, B.C., 89
Cross, G.A.M., see Lamont, G.S., 131
Cushley, W. and Kusel, J.R.
Interaction of the plant toxin ricin with different life cycle stages of *Schistosoma mansoni*, 67

De Jonckheere, J.F.
Characterization of *Naegleria* species by restriction endonuclease digestion of whole-cell DNA, 55
Del Portillo, H.A., Nussenzweig, R.S. and Enea, V.
Circumsporozoite gene of a *Plasmodium falciparum* strain from Thailand, 289
Desjeux, P., see Legrand, D., 117
Donelson, J.E., see Nasir, A., 295
Dore, E., see Pace, T., 193
Dutta, G.P., see Saxena, N., 283

Enea, V., see Del Portillo, H.A., 289
Etges, R., see Bouvier, J., 73
Evans, J.C., see Yarlett, N., 255

Fairlamb, A.H., Henderson, G.B., Bacchi, C.J. and Cerami, A.
In vivo effects of difluoromethylornithine on trypanothione and polyamine levels in bloodstream forms of *Trypanosoma brucei*, 185
Fairlamb, A.H., see Henderson, G.B., 39
Fenton, B.
Peptide digest studies of polymorphic proteins of *Plasmodium falciparum*, 13
Fox, J.A., see Lamont, G.S., 131
Frontali, C., see Pace, T., 193

Garfinkel, L., see Huber, M., 227
Ghatak, S., see Saxena, N., 283
Gibson, S., see Qureshi, F., 137
Gitler, C., see Huber, M., 227

Hedstrom, L., see Verham, R., 1
Henderson, G.B., Fairlamb, A.H. and Cerami, A.
Trypanothione dependent peroxide metabolism in *Crithidia fasciculata* and *Trypanosoma brucei*, 39
Henderson, G.B., see Fairlamb, A.H., 185
Henriquez, D., see Boschetti, M.A., 175
Huber, M., Garfinkel, L., Gitler, C., Mirelman, D., Revel, M. and Rozenblatt, S.
*Entamoeba histolytica*: cloning and characterization of actin cDNA, 227

Iams, K.P., see Shah, J.S., 163

Kennedy, M.W., see Qureshi, F., 137
Killian, H.-D., see Wittich, R.-M., 155
Kimmel, B.E., see Shah, J.S., 163
King, D.L., Chqng, Y.-D. and Turco, S.J.
Cell surface lipophosphoglycan of *Leishmania donovani*, 47
Komuniecki, R., Campbell, T. and Rubin, N.
Anaerobic metabolism in *Ascaris suum*: acyl CoA intermediates in isolated mitochondria synthesizing 2-methyl branched-chain fatty acids, 147
Kusel, J.R., see Cushley, W., 67

Lamont, G.S., Fox, J.A. and Cross, G.A.M.
Glycosyl-*sn*-1,2-dimyristylphosphatidylinositol is the membrane anchor for *Trypanosoma equiperdum* and *T. (Nannomonas) congolense* variant surface glycoproteins, 131
Legrand, D., Desjeux, P., Le Pont, F., Brénière, S.F., Lemesre, J.-L., Santoro, F. and Capron, A.
Identification of a major 72 kilodalton surface antigen in twelve isolates of *Leishmania braziliensis braziliensis*, 117
Lemesre, J.-L., see Legrand, D., 117
Le Pont, F., see Legrand, D., 117
Lloyd, D., see Yarlett, N., 255
Lockwood, B.C., North, M.J., Scott, K.I., Bremner, A.F. and Coombs, G.H.
The use of a highly sensitive electrophoretic method to compare the proteinases of trichomonads, 89

Meek, T.D., see Verham, R., 1
Melendy, T. and Ray, D.S.
Purification and nuclear localization of a type I topoisomerase from *Crithidia fasciculata*, 215
Mercer, J.G., Munn, A.E. and Rees, H.H.
*Echinococcus granulosus*: occurrence of ecdysteroids in protoscoleces and hydatid cyst fluid, 203
Miles, M.A., see Bishop, R.P., 263

Miller, H.R.P., see Qureshi, F., 137
Mirelman, D., see Huber, M., 227
Munn, A.E., see Mercer, J.G., 203

Nasir, A., Cook, G.A. and Donelson, J.E.
Sequences of two kinetoplast minicircle DNAs of *Trypanosoma (Nannomonas) congolense*, 295
Nduka, N. and Okolie, E.E.
Detection of adenosine diphosphate-ribosyl transferase activity in the worm, *Onchocera volvulus*, 97
North, M.J., see Lockwood, B.C., 89
Nussenzweig, R.S., see Del Portillo, H.A., 289

Okolie, E.E., see Nduka, N., 97

Pace, T., Ponzi, M., Dore, E. and Frontali, C.
Telomeric motifs are present in a highly repetitive element in the *Plasmodium berghei* genome, 193
Pandey, V.C., see Saxena, N., 283
Piras, M.M., see Boschetti, M.A., 175
Piras, R., see Boschetti, M.A., 175
Ponzi, M., see Pace, T., 193
Powell, J.W., see Christie, D.A., 125

Qureshi, F., Kennedy, M.W., Gibson, S., Ye, P.Z. and Miller, H.R.P.
Resistance of nematode secretory products to cleavage by mast cell proteinases, 137

Ray, D.S., see Melendy, T., 215
Rees, H.H., see Mercer, J.G., 203
Revel, M., see Huber, M., 227
Robert-Géro, M., see Vedel, M., 81
Rowlands, C.C., see Yarlett, N., 255
Rozenblatt, S., see Huber, M., 227
Rubin, N., see Komuniecki, R., 147

Samaras, N., see Spithill, T.W., 23
Santoro, F., see Legrand, D., 117
Saxena, A., see Saxena, N,, 283
Saxena, N., Saxena, A., Dutta, G.P., Ghatak, S. and Pandey, V.C.
Effect of *Plasmodium yoelii nigeriensis* infection and chloroquine on the hepatic mixed function oxidase system of mice, 283
Scallon, B.J., Bogitsh, B.J. and Carter, C.E.
Cloning of a *Schistosoma japonicum* gene encoding a major immunogen recognized by hyperinfected rabbits, 237
Scott, K.I., see Lockwood, B.C., 89
Shah, J.S., Young, J.R., Kimmel, B.E., Iams, K.P. and Williams, R.O.
The 5′ flanking sequence of a *Trypanosoma brucei* variable surface glycoprotein gene, 163
Spithill, T.W. and Samaras, N.
Genomic organization, chromosomal location and transcription of dispersed and repeated tubulin genes in *Leishmania major*, 23
Stables, J.N., see Christie, D.A., 125

Tielens, A.G.M., Van den Heuvel, J.M. and Van den Bergh, S.G.
Differences in intermediary energy metabolism between juvenile and adult *Fasciola hepatica*, 273
Tielens, A.G.M. and Van den Bergh, S.G.
Glycogen metabolism in *Schistosoma mansoni* worms after their isolation from the host, 247
Turco, S.J., see King, D.L., 47
,
Van den Bergh, S.G., see Tielens, A.G.M., 247, 273
Van den Heuvel, J.M., see Tielens, A.G.M., 273
Vedel, M. and Robert-Géro, M.
Characterization of RNA from *Leishmania tropica* and *Leishmania d. donovani* promastigotes, 81
Verham, R., Meek, T.D., Hedstrom, L. and Wang, C.C.
Purification, characterization, and kinetic analysis of inosine 5′-monophosphate dehydrogenase of *Tritrichomonas foetus*, 1

Walter, R.D., see Wittich, R.-M., 155
Wang, C.C., see Verham, R., 1
Watt, R.A., see Christie, D.A., 125
Williams, R.O., see Shah, J.S., 163
Wittich, R.-M., Kilian, H.-D. and Walter, R.D.
Polyamine metabolism in filarial worms, 155

Yarlett, N., Rowlands, C.C., Evans, J.C., Yarlett, N.C. and Lloyd, D.
Nitroimidazole and oxygen derived radicals detected by electron spin resonance in hydrogenosomal and cytosolic fractions from *Trichmonas vaginalis*, 255
Yarlett, N.C., see Yarlett, N., 255
Ye, P.Z., see Qureshi, F., 137
Young, J.R., see Shah, J.S., 163

# Subject Index


Acetate formation, 273
Actin, 227
Adhesion, Attachment, 175
ADP-ribosyl transferase, 97
Aerobic–anaerobic switch, 273
Anaerobic electron transport, 255
Anaerobic mitochondrion, 147
*Ascaris suum*, 131, 147

Barren region, 163
Branched-chain fatty acid synthesis, 147
*Brugia patei*, 155

Cell surface, 47
Cell surface iodination, 117
Cell surface membrane, 175
Cell-free protein synthesis, 81
Cestoda, 203
Chagas' disease, 175
Chloroquine, 283
Chromatin, 97
Chromosome, 23, 263
Circumsporozoite gene, 289
CoA ester, 147
*Crithidia fasciculata*, 29, 215

Difluoromethylornithine, 185
*Dipetalonema viteae*, 125
*Dirofilaria immitis*, 155
cDNA cloning, 227
DNA sequence, 227, 289

Ecdysteroids, 203
*Echinococcus granulosus*, 203
Ectoenzyme, 73
Electron transport, 255
Energy metabolism, 247, 273
*Entamoeba histolytica*, 227
Excretory/secretory antigen, 131

*Fasciola hepatica*, 273
Ferredoxin, 255
Filariasis, 97
Free radical, 255

Gene cloning, 237
Gene conversion, 163
Gene organization, 23
Genome structure, 193
Genomic library, 237
Glucose breakdown, 273
Glucose metabolism, 125
Glutathione, 185
Glycoconjugate, 47
Glycogen, 247
Glycoprotein, 175

Heat shock proteins, 237
Hepatic monooxygenases, 283
High-performance liquid chromatography, 185
Host–parasite interaction, 175
HPLC, 155
Hydatid cyst fluid, 203

Immunoprecipitation, 117
IMP dehydrogenase, 1
In vitro culture, 247
Infection, 175
Inositol, 131
Integral membrane protein, 73

Karyotype, 263
Kinetoplast DNA, 295
Kinetoplastidae, 81

λgt11 expression vector, 237
Lectin, 175
*Leishmania*, 23
*Leishmania amazonensis*, 73
*Leishmania braziliensis*, 73
*Leishmania braziliensis braziliensis*, 117
*Leishmania d. donovani*, 81
*Leishmania donovani*, 47, 73, 263
*Leishmania enriettii*, 73
*Leishmania infantum*, 73
*Leishmania major*, 73
*Leishmania mexicana*, 73
*Leishmania tropica*, 73, 81
Lipid anchor, 131
Lipophosphoglycan, 47
*Litomosoides carinii*, 155

Malaria, 283
Messenger RNA, 81
MGBG, 155
Minicircle DNA sequence, 295
Mixed function oxidase system, 283
Multigene family, 227
Mycophenolic acid, 1

*Naegleria* DNA, 55
*Naegleria fowleri* epidemiology, 55
*Naegleria* taxonomy, 55
Nematode, 131
*Nippostrongylus brasiliensis*, 131
Nitroimidazole, 255
NMR, 125
Nucleus, 215

*Onchocerca volvulus*, 97, 155
Ordered bi bi, 1
Orthogonal field alternation gel electrophoresis, 263

*Pentatrichomonas hominis*, 89
Peptide digests, 13
Peroxide metabolism, 39
Phosphoenol pyruvate carboxykinase, 125
*Plasmodium berghei*, 193
*Plasmodium falciparum*, 13, 289
*Plasmodium yoelii nigeriensis*, 283
Polyamine, 155, 185
Polymorphic proteins, 13
Protease, 73
Protein acylation, 131
Proteinase, 89, 131
Protoscoleces, 203

Quinolinic acid, 125

Rat mast cell proteinase I, 131
Rat mast cell proteinase II, 131
Receptor–ligand, 175
Repetitive DNA, 193
Replication origin, 295
Ribosomal RNA, 81
Ricin, 67
River blindness, 97
RNA, 23

*Schistosoma japonicum*, 237
*Schistosoma mansoni*, 67, 247
SDS-PAGE, 89
Sequence homology, 295
Spin trap, 255
Succinate, 125
Superoxide, 255
Surface antigen, 117

Telomeric sequences, 193
Topoisomerase, 215
Toxicity, 67
*Trichomitus bactrachorum*, 89
*Trichomonas vaginalis*, 89, 255
*Tritrichomonas foetus*, 1, 89
*Trypanosoma (Nannomonas) congolense*, 131
*Trypanosoma brucei*, 39, 163
*Trypanosoma congolense*, 295
*Trypanosoma cruzi*, 175
*Trypanosoma equiperdum*, 131
*Trypanosoma evansi*, 131
Trypanosome, 185
Trypanothione, 39, 185
Tubulin, 23
Two dimensional electrophoresis, 13

Uncompetitive inhibition, 1

Variant surface glycoprotein, 131
VSG gene, 163